



## Prognostic Factors for Pulmonary Embolism

Submitted by Emmanuel Lemoine on Tue, 12/16/2014 - 10:54

Titre Prognostic Factors for Pulmonary Embolism

Type de publication Article de revue

Auteur Sanchez, Olivier [1], Trinquart, Ludovic [2], Caille, Vincent [3], Couturaud, Francis [4], Pacouret, Gérard [5], Meneveau, Nicolas [6], Verschuren, Franck [7], Roy, Pierre-Marie [8], Parent, Florence [9], Righini, Marc [10], Perrier, Arnaud [11], Lorut, Christine [12], Tardy, Bernard [13], Benoit, Marie-Odile [14], Chatellier, Gilles [15], Meyer, Guy [16]

Editeur American Thoracic Society

Type Article scientifique dans une revue à comité de lecture

Année 2010

Langue Anglais

Date 2010/01/15

Numéro 2

Pagination 168 - 173

Volume 181

Titre de la revue American Journal of Respiratory and Critical Care Medicine

ISSN 1073-449X

Mots-clés echocardiography [17], natriuretic peptides [18], Prognosis [19], pulmonary embolism [20]

Résumé en anglais  
Rationale: The short-term prognosis of pulmonary embolism (PE) depends on hemodynamic status and underlying disease. The prognostic value of right ventricular dysfunction and injury is less well established. Objectives: To evaluate prognostic factors of PE in a multicenter prospective cohort study. Methods: Echocardiography, brain natriuretic peptide (BNP), N-terminal-proBNP and cardiac troponin I measurements were done on admission of 570 consecutive patients with an acute PE. A predictive model was based on independent predictors of 30-day adverse events defined as death, secondary cardiogenic shock, or recurrent venous thromboembolism. Measurements and Main Results: At 30 days, 42 patients (7.4%; 95% confidence interval [CI], 5.5-9.8%) had adverse events. On multivariate analysis, altered mental state (odds ratio [OR] 6.8; 95% confidence interval [CI], 2.0-23.3), shock on admission (OR 2.8; 95% CI, 1.1-7.5), cancer (OR 2.9; 95% CI, 1.2-6.9), BNP (OR 1.3 for an increase of 250 ng/L; 95% CI, 1.1-1.6) and right to left ventricle diameter ratio (OR 1.2 for an increase of 0.1; 95% CI, 1.1-1.4) were associated with 30-days of adverse events. The predictive performance of the model was good (area under receiver operating characteristics curve 0.84 [95% CI, 0.78-0.90]), making it possible to develop a bedside prognostic score. Conclusions: BNP and echocardiography may be useful determinants of the short-term outcome for patients with PE, together with clinical findings. Patients with PE can be stratified according to the initial risk of adverse outcome, using a simple score based on clinical, echocardiographic, and biochemical variables.

URL de la notice <http://okina.univ-angers.fr/publications/ua6459> [21]

DOI 10.1164/rccm.200906-0970OC [22]  
Lien vers le document <http://dx.doi.org/10.1164/rccm.200906-0970OC> [22]

---

## Liens

- [1] [http://okina.univ-angers.fr/publications?f\[author\]=10015](http://okina.univ-angers.fr/publications?f[author]=10015)
- [2] [http://okina.univ-angers.fr/publications?f\[author\]=5369](http://okina.univ-angers.fr/publications?f[author]=5369)
- [3] [http://okina.univ-angers.fr/publications?f\[author\]=10094](http://okina.univ-angers.fr/publications?f[author]=10094)
- [4] [http://okina.univ-angers.fr/publications?f\[author\]=10095](http://okina.univ-angers.fr/publications?f[author]=10095)
- [5] [http://okina.univ-angers.fr/publications?f\[author\]=10096](http://okina.univ-angers.fr/publications?f[author]=10096)
- [6] [http://okina.univ-angers.fr/publications?f\[author\]=10097](http://okina.univ-angers.fr/publications?f[author]=10097)
- [7] [http://okina.univ-angers.fr/publications?f\[author\]=10016](http://okina.univ-angers.fr/publications?f[author]=10016)
- [8] <http://okina.univ-angers.fr/pierremarie.roy/publications>
- [9] [http://okina.univ-angers.fr/publications?f\[author\]=10098](http://okina.univ-angers.fr/publications?f[author]=10098)
- [10] [http://okina.univ-angers.fr/publications?f\[author\]=10020](http://okina.univ-angers.fr/publications?f[author]=10020)
- [11] [http://okina.univ-angers.fr/publications?f\[author\]=10019](http://okina.univ-angers.fr/publications?f[author]=10019)
- [12] [http://okina.univ-angers.fr/publications?f\[author\]=10099](http://okina.univ-angers.fr/publications?f[author]=10099)
- [13] [http://okina.univ-angers.fr/publications?f\[author\]=10100](http://okina.univ-angers.fr/publications?f[author]=10100)
- [14] [http://okina.univ-angers.fr/publications?f\[author\]=10101](http://okina.univ-angers.fr/publications?f[author]=10101)
- [15] [http://okina.univ-angers.fr/publications?f\[author\]=10102](http://okina.univ-angers.fr/publications?f[author]=10102)
- [16] [http://okina.univ-angers.fr/publications?f\[author\]=10018](http://okina.univ-angers.fr/publications?f[author]=10018)
- [17] [http://okina.univ-angers.fr/publications?f\[keyword\]=10595](http://okina.univ-angers.fr/publications?f[keyword]=10595)
- [18] [http://okina.univ-angers.fr/publications?f\[keyword\]=10596](http://okina.univ-angers.fr/publications?f[keyword]=10596)
- [19] [http://okina.univ-angers.fr/publications?f\[keyword\]=1341](http://okina.univ-angers.fr/publications?f[keyword]=1341)
- [20] [http://okina.univ-angers.fr/publications?f\[keyword\]=10587](http://okina.univ-angers.fr/publications?f[keyword]=10587)
- [21] <http://okina.univ-angers.fr/publications/ua6459>
- [22] <http://dx.doi.org/10.1164/rccm.200906-0970OC>

Publié sur *Okina* (<http://okina.univ-angers.fr>)